

Exercise 1

Calc. : ✓

Question B1: Oxygen Intake on a Treadmill

A group of athletes volunteered to have their oxygen intake measured whilst running on a treadmill.

The treadmill allows for adjustments to the incline of the run and to the speed at which the belt of the treadmill rotates.

The treadmill's running power can be adjusted by increasing the speed of the treadmill as well as increasing the incline of the run.

In the table, below, you will find data for the volunteer athletes' intake of oxygen, in litres per minute, for different levels of treadmill running power, in Watts.



Power (Watts)	30	60	90	120	150	180
Oxygen intake (litres/minute)	1.54	2.56	3.45	4.08	4.61	4.93

- a) **Draw** a scatter diagram showing oxygen intake, in litres per minute, on the vertical axis, as a function of power, in Watts, on the horizontal axis.
- Scale of Axes: Represent 10 Watts as 0.5 cm on the horizontal axis and one litre/minute as 1 cm on the vertical axis.

3 marks

The data can be modelled using a linear function  $y = a \cdot x + b$  where  $y$  is the oxygen intake and  $x$  is the power.

- b) Use your calculator to **find** the equation of the line of regression of  $y$  on  $x$ , **giving** the values of  $a$  and  $b$  correct to three decimal places.
- c) Use the line of regression, from part b) to **calculate**, correct to two decimal places, the oxygen intake of an athlete running on a treadmill with a power of 200 Watts.
- If you didn't find values for  $a$  and  $b$  in part b) please use  $a = 0.02$  and  $b = 1.16$ .

2 marks

2 marks


d) **Determine** the value of  $\bar{x}$  and  $\bar{y}$ , **rounding** to two decimal places, as appropriate. **Plot** the point  $(\bar{x}, \bar{y})$  on the scatter diagram. 2 marks

e) **Draw** the regression line on your scatter plot. **Describe** the correlation between oxygen intake and treadmill running power. **Explain** why you described the correlation as you did. 3 marks

Consider the logarithmic model:

$$f(x) = 1.94 \cdot \ln(x) - 5.2$$













This model could also be used to model the bivariate data set Power,  $x$ , and Oxygen Intake,  $y$ . Below is the graph of the logarithmic function  $y = f(x)$ .



f) Use the logarithmic model to **calculate**, correct to two decimal places, the oxygen intake of an athlete when the treadmill's running power is 100 Watts. 2 marks

g) **Calculate**, correct to two decimal places, the value of the derivative of the logarithmic function when the treadmill's running power is 100 Watts. 2 marks

h) **Explain** the meaning of the value of the derivative calculated in part g). 2 marks

<p>An athlete would like to adjust the treadmill's running power to permit an oxygen intake of exactly 3 litres/minute.</p>	
<p>i) Use the logarithmic model to <b>determine</b>, correct one decimal place, the power level at which the treadmill must be set, to permit an oxygen intake of three litres/minute.</p>	2 marks
<p>Both the linear model and the logarithmic model fit well to the given data points. However, when using the models for interpolation or extrapolation, one model falls short i.e., it is not as appropriate as thought.</p>	
<p>j) <b>Select</b> the appropriate word(s) from the list:</p>	2 marks
<p style="text-align: center;"><u>Word Choices</u></p> <p style="text-align: center;">A: Linear</p> <p style="text-align: center;">B: Logarithmic</p> <p style="text-align: center;">C: Interpolation</p> <p style="text-align: center;">D: Extrapolation</p>	
<p>to complete the sentence below.</p>	
<p><b>“The _____ model should not be used for _____.”</b></p>	
<p><b>Write</b> the entire sentence on your exam script. <b>Provide</b> reasoning for your choice of word(s).</p>	
<p>In the group of volunteer athletes, 60% were football players, 30% were cross-country runners and 20% of the athletes did neither of these two sports.</p>	
<p>k) Given that an athlete plays football, <b>calculate</b> the probability that this athlete isn't a cross-country runner.</p>	2 marks
<p>l) Out of the 12 Venn diagrams, numbered one through twelve, given below, <b>choose</b> the Venn diagram where the shaded region, matches the probability that you were asked to calculate in the part k).</p>	1 mark
<div style="display: flex; flex-wrap: wrap; justify-content: space-around;"> <div style="margin: 5px;"> <p>1. </p> </div> <div style="margin: 5px;"> <p>2. </p> </div> <div style="margin: 5px;"> <p>3. </p> </div> <div style="margin: 5px;"> <p>4. </p> </div> <div style="margin: 5px;"> <p>5. </p> </div> <div style="margin: 5px;"> <p>6. </p> </div> <div style="margin: 5px;"> <p>7. </p> </div> <div style="margin: 5px;"> <p>8. </p> </div> <div style="margin: 5px;"> <p>9. </p> </div> <div style="margin: 5px;"> <p>10. </p> </div> <div style="margin: 5px;"> <p>11. </p> </div> <div style="margin: 5px;"> <p>12. </p> </div> </div>	

## Exercise 2

Calc. : ✓

### Question B2: Population Growth in Luxembourg

Luxembourg has experienced steep population growth in recent years.

Back in 2002, Luxembourg had 442,000 inhabitants; twenty years later, this number has increased to 621,000. Note, both population figures are to the nearest thousand.


Luxembourg's population growth can be modelled using the function:

$$N(t) = e^{0.017t+13}$$

where  $N(t)$  is the number of inhabitants in Luxembourg after  $t$ , the time, in years.

January 1<sup>st</sup> 2002, is considered when time,  $t$ , is equal to zero.

The graph of function,  $N(t)$ , is shown below:



- a) Use the model to **calculate** the number of inhabitants in Luxembourg in 2012, **rounding** to the nearest thousand inhabitants.

2 marks

- b) **Show** that the model can be rewritten as:

3 marks

$$N(t) = 442,413 \cdot 1.017^t$$

- c) Use the model in part b) to **find** the yearly growth rate of Luxembourg's population, **giving** your answer as a percentage.

2 marks

The derivative of  $N(t)$  is  $N'(t) = 0.017 \cdot e^{0.017t+13}$ .

- d) Use this derivative to **calculate** the following integral:

3 marks

$$\int_0^{20} N'(t) dt$$

**Round** your answer to nearest whole number. **Explain** what the result means in terms of Luxembourg's population growth.

e) The given model is not flawless; **explain** why this model should not be used in the long run.

1 mark

A second model for population growth in Luxembourg is:

$$N_2(t) = \frac{800,000}{0.8 + 0.96^t}$$

f) Using the second model, **determine** the size of the Luxembourgish population in the long run.


2 marks

A rapid increase in population necessitates new infrastructure, such as, a supply of affordable housing, new hospitals, and schools, as well as, better public transport.

Luxtram, Luxembourg's tram, a part of Luxembourg's public transportation system, started operating in 2018.

On Luxtram's route from "Lux Expo" to "Lycee Bonneveg", the tram makes stops at "Universiteit" and "Coque". On the "Universiteit" to "Coque" segment of its journey, the tram takes 20 seconds to get to its full speed of 19 m/s.

The speed-time diagram, below, shows a typical, Luxtram journey between the stops "Universiteit" and "Coque".



Note,  $t$  is the time in seconds and  $v$  is the speed in metres per second.

g) Use the graph, above, to **calculate** the distance between the stops "Universiteit" and "Coque", **giving** your answer in kilometres.

2 marks

If the tram's full speed were to be 15 m/s, instead of 19 m/s, the speed-time diagram between the stops "Universiteit" and "Coque" would look different to the speed-time diagram shown above. The distance between the stops remains unchanged.

h) **Select**, from the options below, the option which correctly describes the changes to the speed-time diagram if the tram's top speed were to be 15 m/s instead of 19 m/s.

2 marks

**Give** a reason as to why you selected the option you did.

Option	Height of Graph at $v = 15$ m/s Comparison to the Height when $v = 19$ m/s	Width of Graph at $v = 15$ m/s Comparison to the Width when $v = 19$ m/s
A	Lower	No change
B	Lower	Narrower
C	Lower	Wider
D	No change	Narrower
E	No change	Wider

<p>As Luxembourg's public transport is free; the tram is a popular means of transport for many students. Luxtram knows that, on a usual school day, 1,500 passengers take the tram, 35%, of whom, are high school students.</p> <p>Note: Passengers travel independently of each other.</p> <p>Let <math>X</math> be the number of high school students who take the tram, out of the 1,500 passengers, who take the tram on a usual school day.</p>	
i) <b>Explain</b> why $X$ is binomially distributed, <b>stating</b> the parameters of this binomial distribution.	2 marks
j) <b>Calculate</b> the expected value and the standard deviation of the number of high school students taking the tram, on a usual school day, <b>rounding</b> to two decimal places where appropriate.	2 marks
k) <b>Calculate</b> , correct to three decimal places, the probability that, on a usual school day, at most 500 high school students use the tram.	2 marks
l) <b>Determine</b> , on a day which is not a usual school day, the total number of passengers taking the tram, if it were expected that 630 high school students would be taking the tram.	2 marks